Toe Pain

Understanding symptoms, treatment and recovery

Toe Pain

A guide to symptoms, treatment and recovery



Understanding toe pain

Toe pain can affect any part of the toe, including joints, nails, or surrounding soft tissue. It can result from injury, overuse, poor footwear, or underlying medical conditions.

Supporting your recovery

Wear shoes with a wide toe box and good cushioning. A physiotherapist can tailor exercises to the cause of your pain.

Causes and symptoms

Understanding what may be contributing to your pain

Common causes

- Injury or trauma - stubbed, jammed, or broken toes
- Arthritis - inflammation of the toe joints
- Gout - sudden, severe pain usually in the big toe
- Bunions - bony bump at the base of the big toe
- Hammer toe / mallet toe - abnormal toe position causing pain
- Ingrown toenails - nail growing into the skin

Who may be affected?

- Older adults with arthritis or gout
- Athletes or people who exercise regularly
- People wearing tight or ill-fitting shoes
- Those with foot deformities (bunions, hammer toes)

Common symptoms

- Pain in one or more toes
- Swelling, redness, or tenderness
- Stiffness or difficulty moving the toe
- Pain during walking or standing
- Pain at rest (e.g., gout flare-ups)

Important information

- Pain does not improve after 1-2 weeks
- Severe swelling, redness, or deformity
- Difficulty walking or performing daily activities
- Suspected fracture or gout flare-up
- Ingrown toenails with infection

Recovery outlook

Most toe pain improves with conservative care within a few weeks. Early assessment and consistent exercises reduce the risk of chronic pain or deformity.

Exercises and rehabilitation

Gentle movements under professional guidance

Exercise examples

Ask your physiotherapist which exercises are suitable for you. Use controlled movements and work within comfort.



Exercise instructions

1. Toe Stretch: Gently pull toes upward or spread them apart, hold 10-20 seconds
2. Toe Flexion & Extension: Curl and straighten toes 10-15 times
3. Towel Scrunches: Place a towel on the floor and scrunch it with toes 10-15 times

Recovery guidance

Most toe pain improves with conservative care within a few weeks. Early assessment and consistent exercises reduce the risk of chronic pain or deformity.

Management and next steps

Self-care, treatment and when to seek help

Self-care and home management

- Rest - avoid activities that worsen pain
- Ice - 10-15 minutes, 2-3 times daily for injury or swelling
- Footwear - wear shoes with wide toe boxes and good cushioning
- Elevation - raise foot to reduce swelling
- Pain relief - paracetamol or NSAIDs if safe for you
- Avoid tight socks or shoes

Treatment physiotherapists may offer

- Hands-on techniques: mobilization, stretching, massage
- Exercise programs for strengthening and flexibility
- Advice on footwear, orthotics, or taping
- Education on activity modification to reduce pain

Questions or concerns?

- Pain does not improve after 1-2 weeks
- Severe swelling, redness, or deformity
- Difficulty walking or performing daily activities
- Suspected fracture or gout flare-up
- Ingrown toenails with infection

References

- NHS - Toe Pain: www.nhs.uk/conditions/toe-pain
- Chartered Society of Physiotherapy - Foot & Ankle Guidance

B-Well Physiotherapy Services

0300 303 4811

bwell.mskberkshire@nhs.net

www.bwellberkshire.co.uk